

DEPRESSION, DIABETES AND METABOLIC-NUTRITIONAL FACTORS IN ELDERLY HISPANICS

DEPRESSION, DIABETES AND METABOLIC-NUTRITIONAL FACTORS IN ELDERLY HISPANICS

L.J. FITTEN^{1, 2, 3, 5}, F. ORTIZ^{1, 2, 3, 5}, L. FAIRBANKS^{1, 2, 5}, M. ROSENTHAL⁷, G.N. COLE^{1, 2, 5, 7},
F. NOURHASHEMI^{2, 6}, M.A. SANCHEZ^{1, 2}

1. Alzheimer Disease Research Center, David Geffen School of Medicine at UCLA; 2. Greater Los Angeles Veterans Affairs Healthcare System, Sepulveda Campus; 3. Department of Psychiatry Olive View-UCLA Medical Center; 4. Department of Psychiatry and Biobehavioral Sciences, School of Medicine at UCLA; 5. Department of Neurology, David Geffen School of Medicine at UCLA; 6. Purpan University Hospital, Faculty of Medicine, Toulouse, France; 7. Geriatric Research, Education, Clinical Center, Greater Los Angeles Veterans Affairs Healthcare System. This study was supported by funding from the UCLA Alzheimer's Disease Research Center (P50 AG16570); by the State of California Alzheimer's Research and Clinical Center at UCLA, Project No.44137-JC-18796; by the Olive View-UCLA Medical Center, Sylmar, California; and by the VA Greater Los Angeles Healthcare System. Corresponding Author: L. Jaime Fitten, MD, Greater Los Angeles VA Healthcare System, Sepulveda Campus, 16111 Plummer St. (116A-9), Los Angeles, CA. 91343, Telephone: (818) 895-9541, Fax: (818) 895-9515, E-mail: jfitten@ucla.edu. Alternate Corresponding Author: Freddy Ortiz, Ph.D., e-mail: freddyo@ucla.edu

Abstract: *Objective:* To examine the relationship of depression to metabolic and nutritional risk factors in older Hispanics. *Design:* Crosssectional study. *Setting:* Subjects were part of a community-based, cognitive evaluation project that examined 301 subjects in the Eastern San Fernando Valley of Southern California. *Participants:* Two elderly Hispanic groups: 53 clinically depressed, with memory complaints but not demented subjects, and 33 generally healthy, cognitively asymptomatic subjects. *Measurements:* The results of functional and nutritional questionnaires, a medical and neurological examination, 12-hour fasting clinical laboratory tests, MRI or CT scans, and neuropsychological testing. *Results:* Both groups were nearly identical along socio-demographic variables. However, the depressed group differed significantly from the general healthy group not only in percent of diabetics (38% vs.18%), but in the amount of poorly controlled diabetes, and the depressed group consumed about half the amount of fish that the generally healthy group did. *Conclusions:* This study suggests that factors such as poorly controlled diabetes combined with low consumption of foods high in omega-3 fatty acid content such as sea fish may be associated with an increased risk of developing depression in late life. These factors may be socio-economically and culturally influenced and are therefore amenable to modification.

Key words: Depression, metabolic-nutritional factors, elderly hispanics, sea fish.

Introduction

The number of elderly Hispanics in the United States, is expected to grow exponentially in the next three decades nearly tripling from the current 5% of the total U.S. Hispanic population of more than 40 million (1). Hispanics suffer more frequently than non-Hispanics from diabetes, obesity, depression and cognitive impairment (2-5). Depression has been found to be a risk factor for developing cognitive impairment (6), cardiovascular disease (7), and type 2 diabetes (8). Depression is more common among those with diabetes, and depressive symptoms increase a pattern of physiological risk that contributes to the metabolic syndrome (9, 10). Furthermore, high levels of depressive symptoms concomitant with major chronic medical conditions elevate the risk for death among older Mexican Americans (11, 12).

Diabetes is also independently associated with a greater incidence of cognitive impairment (13, 14) with a worse prognosis following these conditions. The co-existence of depressive symptoms and diabetes further extends risk for cognitive decline and functional disability (15). Alzheimer disease is up to four times more common in diabetics (16).

Studies suggest that nutritional factors may play an important role in the development or prevention of diabetes, depression and cognitive impairment or even dementia (17, 18).

For example, diets rich in seafood with high taurine content are thought to help prevent obesity and diabetes (19). Studies also link depression to nutritional changes. Sweet, fatty foods long have been suggested to alleviate stress in vulnerable people through actions that enhance serotonergic activity (20, 21).

We identified a subgroup of subjects with previously undiagnosed depression from an ongoing community based study of cognitive and health status of aged Hispanics. This paper examines the relationship of depression to metabolic and nutritional risk factors, and functional capacity in depressed but non-demented, elderly Hispanic subjects with a cognitive complaint, and compares them to healthier, no complaint, elder Hispanics from the same community. Based on the cited literature, we expected depression to be associated with a higher incidence of type 2 diabetes, more functional impairment, and lower dietary fish consumption.

Methods

Subjects

Subjects were recruited from a large community of elderly Hispanics by lectures on cognitive problems of aging given by research team members at diverse community venues such as local senior centers, churches, primary health care professionals, adult community schools and social services

THE JOURNAL OF NUTRITION, HEALTH & AGING©

agencies (i.e. Meals on Wheels, Adult Day Care Centers, etc.). Other community outreach (recruitment) methods included Spanish radio and television talk shows, bilingual community flyer distributions, and articles published in local Spanish-language newspapers. Subjects were identified and recruited through community outreach efforts conducted by a fully bilingual, bicultural research staff.

Subjects were drawn exclusively from the greater Eastern San Fernando Valley which has an estimated population of approximately one million, with a racial/ethnic composition of 56% Hispanic, 33% White non-Hispanic, 5% African American, 5.4 Asian-Pacific Islander and 0.4% other (22). Two-way transportation service, with door-to-door pick-up was provided for those who did not have the transportation means to come to the medical center for an evaluation.

A total of 301 participants received complete neuropsychiatric evaluations including medical and psychiatric histories, neuropsychological testing, neurological exam, clinical laboratory tests and either CT or MRI scans. Fifty three clinically depressed subjects and a comparison group of 33 generally healthy, asymptomatic elders were identified from the larger group and form the basis of the present study.

The depressed population was defined as outreached, community dwelling Hispanic elders, age 55 or older, with no previous diagnosis of a clearly stated cognitive complaint, but who met DSM IV criteria for clinical depression after neuropsychiatric evaluation. The over 55 years limit was chosen because of similar levels of dysfunction to 65 year old non-Hispanic White populations, as evidenced by published and preliminary studies (23, 24). Table 1 describes the study inclusion and exclusion criteria. The healthy comparison group consisted of 33 identically outreached, community-dwelling Hispanic volunteers aged 55 years or older with no cognitive complaints or mood symptoms and no serious, concurrent, acute or unstable medical conditions, no concurrent non-dementing, central neurologic disease or previous major psychiatric illness but who showed an interest in participating in this memory related research. Relatives of subjects were excluded. Diabetes, hypertension and other features of metabolic syndrome did not form part of the exclusionary criteria for either group. The comparison group was used in this study to evaluate the impact of diabetes and dietary fish intake on mood.

Assessments and diagnoses

This study was approved by the Internal Review Board at Olive View-UCLA Medical Center. Informed consent in verbal and written Spanish or English was obtained from all participants. Consented symptomatic subjects were required to have a designated informant that had daily contact with the subject. All subjects initially received the Mini-Mental Status Examination (25) and the validated Spanish version of the Mini-Mental Status Examination adjusted for age and education (MMSAdj) (26) and questionnaires eliciting demographic, social, family, medical and medication information. The

MMSAdj version (0-30 points) is a screening test for cognitive and memory impairment that has shown better sensitivity and specificity than the traditional MMSE, for detecting cognitive impairment across education levels.

Table 1

Inclusion criteria
1. Hispanic, or of Hispanic origin, background or descendent
2. Aged 55 or over
3. Meets DSM IV criteria for recent minor or major depression
4. Recent history of at least one memory or cognitive complaint
5. English or Spanish speaking
6. Stable health status
Exclusion Criteria
1. Younger than 55 years of age
2. Presence of acute systemic or severe medical illness
3. History or presence of any previous diagnosed major psychiatric illness (e.g. schizophrenia, schizoaffective, bipolar disorder, Obsessive Compulsive Disorder, Major Depression Disorder, etc.)
4. History of psychiatric hospitalization
5. Presence of any central nervous system disease (e.g. dementia, stroke, neurodegenerative disease)
6. Ethnic background other than Hispanic

Hispanics in our sample were identified by self-identification, place of birth, ancestry and language. The examining investigators, all of whom are native Hispanics, conducted the identification procedures. Hispanic subjects' self-reported Spanish and English language proficiency was assessed, as was their level of acculturation to the United States by using the Marin acculturation scale (27). The presence of depression was established by experienced clinicians according to criteria for Major or Minor Depression of the DSM-IV-TR (28). All cases of depression were deemed clinically significant and warranted treatment. The exclusion of subjects with dementia was based on application of criteria for dementia of the DSM-IV-TR. The metric Body Mass Index (BMI) formula was used to calculate BMI index score (29). Metabolic syndrome was established when at least 3 out of 5 of the following conditions were met: 1) Fasting glucose (FBS) ≥ 110 mg/dL; those on anti-diabetic medications with < 110 mg/dL ; 2) Fasting triglycerides ≥ 150 mg/dL and those with triglycerides < 150 mg/dL on medication for high cholesterol; 3) High density lipoprotein (HDL; female ≤ 50 mg/dL and male ≤ 40 mg/dL); 4) High blood pressure (HBP) (resting systolic ≥ 140 mm Hg and / or diastolic ≥ 90 mm Hg, and those with BP $< 140/90$ mm Hg on anti-hypertensive medications); 5) BMI ≥ 28 kg/m².

All study subjects and comparison subjects received an extensive neuropsychiatric diagnostic assessment before they were stratified into diagnostic groups. The neuropsychiatric evaluation included neuropsychological testing, a neurological examination, twelve-hour fasting laboratory tests (subjects were reminded by phone the night before and asked about fasting in the morning) including, thyroid function tests (TFTs) (TSH, Tri-iodothyronine (T³) and Free thyroxine), vitamin B¹² and folate, fasting blood sugar (FBS), hemogram, kidney panel, lipid panel, liver panel, vitamins A C and E , homocysteine and apolipoprotein (APOE) status, and either brain magnetic resonance imaging (MRI) or computed tomography (CT)

DEPRESSION, DIABETES AND METABOLIC-NUTRITIONAL FACTORS IN ELDERLY HISPANICS

depending on the presence or absence of MRI contra-indications. Additional clinical values obtained included blood pressure (mm/hg), weight (kg) and height (cm). The Geriatric Depression Scale-15 items (GDS) (30) was used to estimate the severity of subjects' depressive symptoms in subjects diagnosed with clinical depression and to confirm the absence of significant depressive symptoms in the comparison group. The normed and standardized neuropsychological tests for Hispanics (31) used in the study included the assessment of attention-concentration, memory, language, visual-spatial, motor coordination, and executive functions. The Spanish version of UCLA Alzheimer Disease Center's Activity of Daily Living Inventory (ADLI) was also administered. A self-reported Seafood Consumption Questionnaire in Spanish was developed for this study by the study's native speakers. The self-reported questionnaire was designed to assess the type of seafood consumed, its origin, amount and frequency of consumption. The amount consumed on a weekly and monthly basis by weight was estimated through the use of three dimensional models of known portion sizes of fish shown directly to subjects. In view of the low fish consumption of many subjects, all data was converted to grams of fish consumed per month. The questionnaire was pilot tested with a focus group of elderly Hispanic subjects and adjusted according to the group feedback on terminology and other facets of the questionnaire and its presentation. Then, the revised questionnaire was retested for validity and accuracy in a small number of elderly Hispanics from the original population (unpublished data). The same native Spanish-speaking investigator presented the Seafood Consumption Questionnaire to all subjects. All diagnoses made in this study were established after diagnostic consensus conferences involving clinicians from the other UCLA ADRC sites.

Data analysis

Differences between depressed and comparison subjects on the demographic and outcome measures described above were tested using two-tailed t-tests for independent groups for continuous variables and the chi square test of independence for categorical variables, with alpha = 0.05. Effects of confounding or correlated variables were controlled by including them as covariates in a univariate analysis of variance with group as a between subject factor.

Results

Subject Characteristics

As shown in table 2 the depressed and comparison groups were very similar along demographic and psychosocial variables, except for differences in age, GDS score and MMSAdj score. The healthier, non-depressed comparison group was significantly older (72.3, ±6.3) than the depressed group (63.7, ±7.0), (p <.000). As expected, the depressed group had significantly higher GDS (10.2, ±3.3), and lower MMSAdj scores (28.2 ±2.5) than the comparison group, GDS (.50, ±1.0)

p < .000, and MMSAdj (29.7, ±0.7) p < .002. No significant differences in education, annual income (estimated average), Marin Acculturation Scale score, weight, height and BMI were found between the groups. The percentage of females was high in both depressed (81.1%) and comparison (84.8%) groups, but no difference between them was found.

Table 2
Psychosocial and Demographic Variables

Demographic and Psychosocial Variables	Groups		Test value	P value
	(n=53) Depression x (SD)	(n=33) Controls x (SD)		
Age	63.7 (7.0)	72.3 (6.9)	F=.018	.000
Gender (% Female)	81.1%	84.8%	-	ns
GDS score ^a	10.2 (3.3)	.50 (1.0)	F= 4.726	.000
MMSAdj score ^b	28.2 (2.5)	29.7 (0.76)	F=22.219	.002
Education (yrs)	6.6 (3.6)	7.9 (3.7)	-	ns
Approximated Annual Mean Income (US\$)	11,084 (7,867)	11,287 (8,752)	-	ns
Marin Acculturation Scale score	1.51 (0.93)	1.58 (1.03)	-	ns
Weight (kg)	68.3 (12.3)	68.8 (12.6)	-	ns
Height (cm)	155.7 (9.7)	154.2 (11.2)	-	ns
BMI (kg/m ²) ^c	28.9 (5.3)	29.0 (4.4)	-	ns

a. Geriatric Depression Scale; b. Mini-Mental Status Examination adjusted for education and Age; c. Body Mass Index

Depression, Type 2 Diabetes and Metabolic Syndrome

The rate of diabetes in the depressed group (38%) was more than twice that of the comparison group (18%), even though mean BMI for both groups was nearly identical. No difference between groups was found in pharmacologic treatment for diabetes. Two thirds of diabetics in each group received treatment. Non-treated diabetics either did not know they had diabetes or chose not to take treatment. Figure 1 shows that despite reported treatment, a significant difference in mean fasting glucose level was found between the depressed (124.3, ±71.3), and the comparison (94.2, ±21.2), p < .000) groups. A univariate analysis of variance, using BMI as a covariate, found that the effect of depression on fasting glucose levels was independent of BMI (F=4.899, p < .030) in these groups.

Table 3 shows the list of outcome variables. As shown, statistically significant differences were found between the depressed and the control groups on mean fasting glucose (elevated in the depressed group, 124.3, ±71.3 vs. 94.2 ±21.2; p < .021), ADLI mean scores (higher in the depressed group, 11.5 ±4.8 vs. 8.1 ±0.5; p < .000), and mean grams of fish consumption per month (lower in the depressed group, 134.7, ±19.0 vs. 162.5, ±58.4; p < .000). However, a significantly lower systolic blood pressure was found for the depressed subjects (134.7, ±19.0 vs. 148.7, ±26.0; p < .006). Vitamin levels were normal in both groups and no other statistically significant differences were found in measured diabetic and vascular disease risk factors.

Table 3
Groups Outcome Variables

Psychosocial, Nutritional and Metabolic Variables	Normal ranges	Groups		Test Value (F)	P value*
		Depression (n=53) mean (SD)	Controls (n=33) Mean (SD)		
Fasting glucose (mg/dL)	60-109	124.3 (71.3)	94.2 (21.2)	5.516	.021
ADLI score†	-	11.5 (4.8)	8.1 (.5)	14.540	.000
Fish consumption (gm/month)	-	89.2 (80.7)	162.5 (58.4)	19.169	.000
Systolic blood pressure (mm/Hg)	90-120	134.7 (19.0)	148.7 (26.0)	7.958	.006
Diastolic blood pressure (mm/Hg)	60-80	76.1 (11.0)	73.9 (13.5)	-	ns
Total cholesterol (mg/dL)	< 199	210.3 (35.4)	209.6 (39.0)	-	ns
Triglycerides (mg/dL)	< 149	176.8 (76.2)	210.6 (162.4)	-	ns
Low density lipoprotein (mg/dL)	< 100	118.4 (37.3)	117.9 (33.0)	-	ns
High density lipoprotein (mg/dL)	> 40	56.5 (16.1)	52.9 (16.3)	-	ns
Homocysteine (mmol/l)	< 10.3	9.5 (3.7)	10.7 (2.0)	-	ns
Metabolic syndrome (%)		56%	55%	-	ns

* Analysis of Variance; † Activity of Daily Living Inventory

No statistically significant difference was found between groups on the rate of diabetes plus pre-diabetic metabolic syndrome (54.7% in the depressed group vs. 54.5% in the control group). The comparison group had a higher rate of pre-diabetic metabolic syndrome (44.4%) than the depressed group (27.3%) after controlling for diabetes, but the difference was not statistically significant. Certain subjects in both groups who had metabolic syndrome were treated: Thirty percent (10/33) of comparison subjects were on cholesterol lowering agents and 45% (15/33) on anti-hypertensive medications, while in the depressed group, 26% (14/53) reported being on cholesterol-lowering drugs and 42% (22/53) on anti-hypertensive medication. No specific triglyceride therapy was reported by either group. Cholesterol levels did not correlate with level of depression and no group differences were found in possible depressogenic medications (e.g. propranolol, corticosteroids, etc.) or drugs known to increase insulin resistance.

Depression and Daily Functioning

Measures in Activity of Daily Living Inventory (ADLI) between the depressed (11.5, ±4.8) and comparison group (8.1, ±0.5) differed significantly (p < .000), with the depressed group scores showing more functional impairment than the comparison group (See Table 2). This difference held after including MMSAdj as a covariate in the analysis, (F=6.625, p. 012), indicating that the group difference was related to depression and not to the small difference between groups in cognitive status (28.2 vs. 29.7).

Depression and Diet

Whereas several depressed subjects reported diminished enjoyment of food, substantial alterations of eating patterns and major weight loss were not reported. Analysis of weight, BMI and vitamin values between depressed and comparison subjects revealed no significant differences (Table 2). However, analysis of estimated fish consumption through the Seafood Consumption Questionnaire (SFQ) yielded notable results. There was an overwhelming preference and consumption in both groups for salt water over lake or stream fish. Figure 1 illustrates monthly mean fish consumption for the comparison group (162.5, ±58.4) versus the depressed group (89.2, ±80.7, p. < .000). A univariate analysis of variance, using fasting glucose level as a covariate, found that the difference between depressed subjects and controls in fish consumption was independent of fasting glucose level (F=17.128, p < 000) in these groups. This indicates that the effect of fish consumption on depression was not due to dietary effects on diabetes.

Figure 1
All subjects





Figure 2

Mean of fasting glucose level compared between groups



DEPRESSION, DIABETES AND METABOLIC-NUTRITIONAL FACTORS IN ELDERLY HISPANICS

Discussion

This study compared a group of community outreach Hispanic elders who initially voiced a cognitive complaint, but who were found to be depressed and not demented, to a similarly community-recruited group of generally healthy Hispanic elders with no complaint. Originating in the same community, both groups were nearly identical along the psychosocial-demographic continuum. As would be expected the depressed group had higher GDS and ADLI scores, and slightly lower, but still normal, MMSAdj scores than the comparison group. Somewhat unexpected was the older age of the comparison group.

When the groups were compared along multiple metabolic/nutritional variables they were indistinguishable, with the exception of fasting blood glucose level and reported monthly fish consumption where the differences between the depressed and comparison groups were highly significant. The comparison group consumed nearly twice the amount of sea fish as the depressed group and had a group mean fasting blood glucose well within normal limits, whereas the depressed group's mean was almost in the diabetic range. The depressed group had many more diabetic subjects and higher representation of diabetics with fasting blood sugars in the diabetic range than did the comparison group. Furthermore, the comparison group consumed nearly twice the amount of fish per month than did the depressed subjects, and this was an independent effect from diabetic status since the strong difference between groups persisted after diabetics were removed from both groups.

Our results are consistent with previous reports in the literature on the high prevalence of diabetes in Hispanic-American populations, though less is known about diabetes in older Hispanics. Some studies on Hispanics have suggested that the vulnerability to diabetes is related to the amount of American-native admixture present. Both, comparison and depressed groups in this study show a considerably higher prevalence of diabetes than seen in older Americans of European descent even when recent increases in diabetes in the American population at large are taken into consideration. Both study groups' high adiposity is also noteworthy. Striking, however, is the particularly high proportion of diabetics in the somewhat younger depressed group. In all of the depressed subjects the initial complaint was a cognitive one, even though no significant cognitive deficit was found after extensive neuropsychological testing. The literature also reports a relationship between diabetes and subsequent development of depression or eventual development of diseases associated with cognitive decline (8, 32-36). This study is consistent with these reports and in addition shows a strong association between depression and poorer fasting glycemic control.

While the present study design does not permit the determination of causality, it does suggest that tighter glycemic control might help in reducing the risk of developing

symptomatic depression if diabetes is present, although other factors may also be involved. For example, there is a linkage between insulin resistance, the generation of reactive oxygen species and chronic inflammation (37). Lower socio-economic status has been related to greater chronic inflammation including higher tumor necrosis factor (TNF)-alpha in Hispanics (38). Future studies should target the role of good glycemic control on the manifestation of depressive symptoms in older persons with type II Diabetes and on the emergence of treatment resistant depression.

Another finding was the diabetes-independent relationship of amount of sea-fish consumption to depression. While both comparison and depressed groups were very low consumers of fish when compared to consumption amounts advocated by the Food and Drug Administration (12 oz varied sea fish per week), sea-fish consumption in the depressed group was dramatically low (3 oz per month). Sea-fish such as salmon, mackerel, tuna, sardines, etc., are major sources of omega-3 fatty acids (eicosapentanoic acid-EPA, and docosahexanoic acid-DHA) in the diet. Epidemiological, experimental and new clinical studies have shown a strong connection between low consumption of foods rich in omega-3 fatty acids and increased depression (39), though not every study has shown a clear association (40). Animal studies have shown that deficiencies in the omega-3 fatty acids (EPA and DHA) can produce deficits in dendritic arborization and synaptic function, involving neurotransmitters (acetylcholine, serotonin, dopamine) implicated in mood disorders (40, 41) and deficits in BDNF, a major neurotrophic factor with depression associated deficits (42). Supplementation with omega-3 fatty acids typically mitigates the synaptic and neurotransmitter deficits and may also improve arborization, while fish oil supplementation can mitigate depression-related behavior in animal models (43).

NIH studies have shown that higher national fish consumption for a country correlates with lower depression rates when compared to countries consuming the least amount of fish (44, 45). Researchers are now observing increasing rates of depression in regions of the world that are moving away from traditional omega-3 rich diets to typical Western foods with high omega-6 fatty acids intake (39). In a recent clinicopathologic study, McNamara et al. (2006) demonstrated selective deficits in omega-3 fatty acid, DHA, content in the postmortem cortex of patients with Major Depressive Disorder (55). Clinical studies have demonstrated that those with depression do have lower levels of omega-3 fatty acids in the blood (17, 46, 47). Results of clinical trials have shown improvement of depressive symptoms in depressive illness or treatment resistant depression over placebo when subjects were given omega 3 fatty acid treatment in modest doses (48, 49). The mechanisms by which omega-3 fatty acids may influence mood have not been elucidated, but it is clear that these substances are important to brain micro-structure and neurotransmitter metabolism and may have the capacity to lower levels of immune agents such as tumor necrosis factor

alpha (TNF α), interleukin 1 beta (IL-1B), and prostaglandin E2, substances which have been reported to be elevated in depression (17).

Factors which limit the generalizability of the study results include the modest sample size and non-random selection of subjects and the high proportion of women, although the community sample used in this study has advantages over the more traditional samples taken from hospital or outpatient populations. The study is also cross-sectional and correlational in design not allowing for the identification of exclusive causal factors in the development of depression. Finally, the depressed subjects who responded initially to recruitment with a cognitive concern may not be characteristic of all depressed older Hispanics. Furthermore, California Hispanics are overwhelmingly of Mexican and Central American origin and represent about two-third of the Hispanics in the U.S., but are not representation of all U.S. Hispanics many of whom come from other cultural and racially distinct areas.

In conclusion, depressed older Hispanics initially presented with a cognitive complaint but were not demented. The depressed group differed significantly from the healthier, asymptomatic comparison group particularly in the amount of poorly controlled diabetes found. In addition, the depressed group consumed very small quantities of fish, half the amount the comparison group did. This study suggests that factors such as poorly controlled diabetes combined with very low consumption of omega-3 fatty acids may greatly increase the risk of developing depression in late life. These factors may be socio-economically and culturally influenced and are therefore amenable to modification.

All contributors are listed authors and are Federal, VA employees except Dr. F. Nourhashemi who is not a VA employee.

Author Contributions: Indicate authors' role in study (1) concept and design, (2) acquisition of subjects and/or (3) data, analysis and interpretation of data, and (4) preparation of manuscript. (List all author(s) name). Author Name: L. Jaime Fitten, M.D., 1,3,4; Author Name: Freddy Ortiz, M.A., 2,3,4; Author Name: Lynn Fairbanks, Ph.D., 3,4; Author Name: Mark Rosenthal, M.D., 3,4; Author Name: Greg Cole, Ph.D., 4; Author Name: Fatemeh Nourhashemi, M.D., 2; Author Name: Mary Anne Sanchez, 2

Sponsor's Role: Indicate sponsor's role in the design, methods, subject recruitment, data collections, analysis and preparation of paper. NONE

Financial Disclosure(s): Author Name: L. Jaime Fitten, M.D., Speakers Forums-Novartis, Forest, Pfizer; Author Name: Freddy Ortiz, M.A., None; Author Name: Lynn Fairbanks, Ph.D., None; Author Name: Mark Rosenthal, M.D., None; Author Name: Greg Cole, Ph.D., None; Author Name: Fatemeh Nourhashemi, M.D., None; Author Name: Mary Anne Sanchez, None

References

- Administration on Aging. (U.S. Department of Health and Human Services). A Profile of Older Americans: 2003. 2003.
- Dunlop DD, Song J, Lyons JS, Manheim LM, Chang RW. Racial/ethnic differences in rates of depression among pre-retirement adults. *Am J Public Health.* 2003;93(11):1945-52.
- Center for Disease Control. Diabetes among Hispanics--Los Angeles County, California, 2002-2003. *MMWR Morb Mortal Wkly Rep.* 2003;52(47):1152-5.
- Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med.* 2003;163(4):427-36.
- Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA.* 2002;287(3):356-9.
- Sherita Hill Golden JEW, Daniel E. Ford, Hsin-Chieh Yen, Catherine Paton Sanford, F. Javier Nieto, Frederic L Brancati. Depressive symptoms and the risk of type 2 diabetes. *Diabetes Care.* 2004;27(2):429-435.
- Abramson J, Berger A, Krumholz HM, Vaccarino V. Depression and risk of heart failure among older persons with isolated systolic hypertension. *Arch Intern Med.* 2001;161(14):1725-30.
- Golden SH, Williams JE, Ford DE, et al. Depressive symptoms and the risk of type 2 diabetes: the Atherosclerosis Risk in Communities study. *Diabetes Care.* 2004;27(2):429-35.
- Miyata S, Hirano S, Kamei J. Diabetes attenuates the antidepressant-like effect mediated by the activation of 5-HT1A receptor in the mouse tail suspension test. *Neuropsychopharmacology.* 2004;29(3):461-9.
- Ahlberg AC, Ljung T, Rosmond R, et al. Depression and anxiety symptoms in relation to anthropometry and metabolism in men. *Psychiatry Res.* 2002;112(2):101-10.
- Black SA, Markides KS. Depressive symptoms and mortality in older Mexican Americans. *Ann Epidemiol.* 1999;9(1):45-52.
- Rodriguez-Saldana J, Morley JE, Reynoso MT, et al. Diabetes mellitus in a subgroup of older Mexicans: prevalence, association with cardiovascular risk factors, functional and cognitive impairment, and mortality. *J Am Geriatr Soc.* 2002;50(1):111-6.
- Wu JH, Haan MN, Liang J, Ghosh D, Gonzalez HM, Herman WH. Impact of diabetes on cognitive function among older Latinos: a population-based cohort study. *J Clin Epidemiol.* 2003;56(7):686-93.
- Raji MA, Al Snih S, Ray LA, Patel KV, Markides KS. Cognitive status and incident disability in older Mexican Americans: findings from the Hispanic established population for the epidemiological study of the elderly. *Ethn Dis.* 2004;14(1):26-31.
- Nguyen HT, Black SA, Ray LA, Espino DV, Markides KS. Predictors of decline in MMSE scores among older Mexican Americans. *J Gerontol A Biol Sci Med Sci.* 2002;57(3):M181-5.
- Zhao L, Teter B, Morihara T, et al. Insulin-degrading enzyme as a downstream target of insulin receptor signaling cascade: implications for Alzheimer's disease intervention. *J Neurosci.* 2004;24(49):1120-6.
- Bourre JM. Roles of unsaturated fatty acids (especially omega-3 fatty acids) in the brain at various ages and during ageing. *J Nutr Health Aging.* 2004;8(3):163-74.
- Gonzalez-Gross M, Marcos A, Pietrzik K. Nutrition and cognitive impairment in the elderly. *Br J Nutr.* 2001;86(3):313-21.
- Tsuboyama-Kasaoka N, Shozawa C, Sano K, et al. Taurine (2-aminoethanesulfonic acid) deficiency creates a vicious circle promoting obesity. *Endocrinology.* 2006;147(7):3276-84.
- Leigh Gibson E. Emotional influences on food choice: Sensory, physiological and psychological pathways. *Physiol Behav.* 2006.
- Ford ES, Mokdad AH, Giles WH, Brown DW. The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. *Diabetes.* 2003;52(9):2346-52.
- San Fernando Valley Economic Research Center. (College of Business and Economics, California State University of Northridge). San Fernando Valley Economic Report 2004-2005. 2004 October 2004.
- Clark CM, DeCarli C, Mungas D, et al. Earlier onset of Alzheimer disease symptoms in Latino individuals compared with Anglo individuals. *Arch Neurol.* 2005;62(5):774-8.
- Fitten LJ, and Ortiz, F. Early onset of dementia in a community outreach sample of elderly Hispanics. Sylmar, CA; 2006.
- Folstein MF, Folstein, S.E., & McHugh, P.R. Mini-Mental State: a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research.* 1975;12:185-198.
- Mungas D, Marshall SC, Weldon M, Haan M, Reed BR. Age and education correction on mini-mental state examination for English- and Spanish-speaking elderly. *Neurol.* 1996;46:700-06.
- Marin G, Sabogal F, Marin BV, Otero-Sabogal R, Pérez-Stable EJ. Development of a short acculturation scale for Hispanics. *Hispanic Journal of Behavioral Science.* 1987;9:183-205.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders, Fourth Edition, Text Revision (DSM-IV-TR). 4th ed Washington, D. C.: American Psychiatric Association; 2000.
- Keys A, Fidanza F, Karvonen MJ, Kimura N, Taylor HL. Indices of relative weight and obesity. *J Chronic Dis.* 1972;25(6):329-43.
- Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. *Clinical Gerontologist.* 1986;5(1-2):165-173.
- Pontón MO, Satz P, Herrera L, et al. Normative data stratified by age and education for the Neuropsychological Screening Battery for Hispanics (NeSBHIS) : A standardization report. *J Inter Neuropsych Soc.* 1996;2:96-104.
- Thomas AJ, Kalaria RN, O'Brien JT. Depression and vascular disease: what is the relationship? *J Affect Disord.* 2004;79(1-3):81-95.

DEPRESSION, DIABETES AND METABOLIC-NUTRITIONAL FACTORS IN ELDERLY HISPANICS

33. McKellar JD, Humphreys K, Piette JD. Depression increases diabetes symptoms by complicating patients' self-care adherence. *Diabetes Educ.* 2004;30(3):485-92.
34. Egede LE. Diabetes, major depression, and functional disability among U.S. adults. *Diabetes Care.* 2004;27(2):421-8.
35. McCaffery JM, Niaura R, Todaro JF, Swan GE, Carmelli D. Depressive symptoms and metabolic risk in adult male twins enrolled in the National Heart, Lung, and Blood Institute twin study. *Psychosom Med.* 2003;65(3):490-7.
36. Saydah SH, Brancati FL, Golden SH, Fradkin J, Harris MI. Depressive symptoms and the risk of type 2 diabetes mellitus in a US sample. *Diabetes Metab Res Rev.* 2003;19(3):202-8.
37. Melek C, Arkan ALH, Florian R, Greten, Shin Maedan, Zhi-Wei Li, Jeffrey M. Long, Anthony Wynshaw-Boris, Giuseppe Poli, Jerrold Olefsky, and Michael Karin. IKK- β links inflammation to obesity-induced insulin resistance. *Nature medicine.* 2005;11(2):191-198.
38. Annemarie Koster. HB, Brenda W. J. H. Penninx, Anne B. Newman, Tamara B. Harris, Jacques Th. M. van Eijk, Gertrudis I. J. M. Kempen, Eleanor M. Simonsick, Karen C. Johnson, Ronica N. Rooks, Hilsa N. Ayonayon, Susan M. Rubin, and Stephen B. Kritchevsky. Association of inflammatory markers with socioeconomic status. *J Gerontol: Med Sc.* 2006;61A(3):284-290.
39. Sontrop J, Campbell MK. Omega-3 polyunsaturated fatty acids and depression: a review of the evidence and a methodological critique. *Prev Med.* 2006;42(1):4-13.
40. Parker G, Gibson NA, Brotchie H, Heruc G, Rees AM, Hadzi-Pavlovic D. Omega-3 fatty acids and mood disorders. *Am J Psychiatry.* 2006;163(6):969-78.
41. McNamara RK, Ostrander M, Abplanalp W, Richtand NM, Benoit SC, Clegg DJ. Modulation of phosphoinositide-protein kinase C signal transduction by omega-3 fatty acids: Implications for the pathophysiology and treatment of recurrent neuropsychiatric illness. *Prostaglandins Leukot Essent Fatty Acids.* 2006.
42. Rao JS, Ertley RN, Lee HJ, et al. n-3 Polyunsaturated fatty acid deprivation in rats decreases frontal cortex BDNF via a p38 MAPK-dependent mechanism. *Mol Psychiatry.* 2006.
43. Naliwaiko K, Araujo RL, da Fonseca RV, et al. Effects of fish oil on the central nervous system: a new potential antidepressant? *Nutr Neurosci.* 2004;7(2):91-9.
44. Jacka EN, Pasco JA, Henry MJ, Kotowicz MA, Nicholson GC, Berk M. Dietary omega-3 fatty acids and depression in a community sample. *Nutr Neurosci.* 2004;7(2):101-6.
45. Bourre JM. [Omega-3 fatty acids in psychiatry]. *Med Sci (Paris).* 2005;21(2):216-21.
46. Maes M, Christophe A, Delanghe J, Altamura C, Neels H, Meltzer HY. Lowered omega3 polyunsaturated fatty acids in serum phospholipids and cholesteryl esters of depressed patients. *Psychiatry Res.* 1999;85(3):275-91.
47. Su KP, Shen WW, Huang SY. Are omega3 fatty acids beneficial in depression but not mania? *Arch Gen Psychiatry.* 2000;57(7):716-7.
48. Gao K, Calabrese JR. Newer treatment studies for bipolar depression. *Bipolar Disord.* 2005;7 Suppl 5:13-23.
49. Frangou S, Lewis M, McCrone P. Efficacy of ethyl-eicosapentaenoic acid in bipolar depression: randomized double-blind placebo-controlled study. *Br J Psychiatry.* 2006;188:46-50.

**XIXth IAGG World Congress of
GERONTOLOGY
and GERIATRICS**

LONGEVITY, HEALTH AND WEALTH

IAGG
5-9 JULY 2009
PALAIS DES
CONGRÈS
PARIS

Mark
your
calendar!

Abstract submission deadline: 31 January 2009
www.gerontologyparis2009.com